

# Carbon Flux related to Peatland Use Change

Wiwiek Setyawati, Waluyo Eko cahyono, Rosida, Saipul Hamdi, Atep Radiana  
The Centre For Atmospheric Science and Technology – LAPAN Indonesia

## SUMMARY


- CO<sub>2</sub> flux over Sumatra is relatively higher than over Kalimantan and Papua peatlands. It is probably due to different degree of peatland cover over that influence the soil water content and surface temperature. Further study relating carbon flux and peatland cover probably by using NDVI data is needed.
- Hydrological systems over most peatland in Sumatra and Southern Papua were governed by precipitation rate, probably it is the only water input. But in most peatland area in Kalimantan and little part of Papua, there are other factors that might control the peatland hydrological system. Therefore, further study is needed to investigate the cause of indifferences.

## INTRODUCTION


- Conversion of peatland to agricultural or plantation land have resulted in changes in the hydrological system, such as change in water level and also the peat temperature and humidity.
- Change in biological conditions from anaerobes to aerobes results in an increase in the respiration activity of microorganisms on peat land which results in an increase of CO<sub>2</sub> (carbon dioxide) emissions into the atmosphere.
- Precipitation as one of water input to the peatland hydrology might also influence peat water table thus carbon emission
- The study aimed to determine carbon flux from various peat water table and to find relationship between precipitation and peat water table.

## METHODE

- The study focused on peatland over Sumatra, Kalimantan and Papua, Indonesia




- Data processing were shown in the flowchart.



- The equation used to calculate peat water table and carbon flux (Astiani et al., 2016):

$$\text{Soil } T \text{ in } ^\circ\text{C} = 35,99 - (0,105 \times \text{peat water table in cm})$$


$$\begin{aligned} & \sqrt[2]{\text{flux CO}_2 \text{ in } \mu\text{mol}/\text{m}^2\text{s}^1} \\ &= 1,756 + (0,0297 \times \text{peat water table}) - (0,0204 \times \text{soil water content}) \\ &+ (0,0220 \times \text{surface } T) \end{aligned}$$



A graphic used to convert water moisture unit from kg/m<sup>2</sup> to water content in % wet basis for top 10 cm soil by assuming peat bulk density 0.1 g/cm<sup>3</sup>


## PEATLAND CO<sub>2</sub> FLUX

- CO<sub>2</sub> flux over Sumatra is relatively higher than over Kalimantan and Papua peatlands.
- It is probably due to less peatland cover over Sumatra than Kalimantan and Papua that influence the soil temperature and water content and also surface temperature.



## PEAT WATER TABLE AND PRECIPITATION RATE

- Peat water tables in the North-eastern Sumatra, Southern Papua, and northern and eastern Kalimantan are inversely proportional to precipitation rates. On the contrary, peat water table in the western and southern Kalimantan is directly proportional to precipitation rate.
- It can be concluded that hydrological systems over most peatland in Sumatra and Southern Papua were governed by precipitation rate, probably the only water input. But in most peatland area in Kalimantan and little in Papua, there might be other factors that control the peatland hydrological system



## FURTHER STUDY

- Further study to investigate relation between CO<sub>2</sub> flux and peatland cover probably by using data NDVI, and also factors that control hydrological system in Kalimantan and little part of Papua are required.
- In situ measurements of Soil T and water content, surface T, Precipitation and peat water table are needed to validate the input and output of the model used in this study.

## ACKNOWLEDGEMENT

- We thanks to Giovanni NASA for allowing us to use the data freely. This work was funded by the Centre for Atmospheric Science and Technology – LAPAN and Kemenristekdikti through INSINAS programme

## REFERENCES

- Couwenberg et al., 2010. Greenhouse gas fluxes from tropical peatlands in South-east Asia. Global Change Biology, 16, 1715 – 1732.
- Astiani et al., 2016. Effects of water table level on soil CO<sub>2</sub> respiration in West Kalimantan forested and bare peatland: An experimental stage. Nusantara Bioscience, 8, No. 2, pp. 201-206.
- Mezhabuddin et al., 2014. Modelling effects of seasonal variation on water table depth on net ecosystem CO<sub>2</sub> exchange of tropical peatland. Biogeosciences, 11, pp. 577-599.
- Silvola et al., 1996. CO<sub>2</sub> fluxes from peat in boreal mires under varying temperature and moisture conditions. J Ecol, 84, pp. 219-228.
- Chimner et al., 2003. Influence of water table position on CO<sub>2</sub> emissions in a Colorado subalpine fen: an in situ microcosm study. Soil Biol and Biogeochem, 35, pp. 345-351.
- NASA; 2018. Giovanni The Bridge Between Data and Science v 4.30; <https://giovanni.gsfc.nasa.gov/giovanni/>



Wiwiek Setyawati  
[wiwieksetyawati21@gmail.com](mailto:wiwieksetyawati21@gmail.com);  
[wiwiek.setyawati@lapan.go.id](mailto:wiwiek.setyawati@lapan.go.id)